

RayBio® Mouse Atherosclerosis Antibody Array 1

Detect 22 Atherosclerosis factors in one experiment

| | A | B | C | D | E | F | G | H |
|---|--------|--------------|--------------|---------------|--------------|-------|----------------|------------|
| 1 | POS | POS | NEG | NEG | Blank | bFGF | CD40 | Eotaxin |
| 2 | POS | POS | NEG | NEG | Blank | bFGF | CD40 | Eotaxin |
| 3 | GCSF | GM-CSF | IFN γ | IL-1 α | IL-1 β | IL-2 | IL-3 | IL-4 |
| 4 | GCSF | GM-CSF | IFN γ | IL-1 α | IL-1 β | IL-2 | IL-3 | IL-4 |
| 5 | IL-5 | IL-6 | IL-13 | L-Selectin | MCP1 | M-CSF | MIP-3 α | P-Selectin |
| 6 | IL-5 | IL-6 | IL-13 | L-Selectin | MCP1 | M-CSF | MIP-3 α | P-Selectin |
| 7 | RANTES | TNF α | VEGF | Blank | Blank | Blank | Blank | POS |
| 8 | RANTES | TNF α | VEGF | Blank | Blank | Blank | Blank | POS |

* For use with serum, plasma, condition medium, urine and all other body fluids, cell lysates and certain tissue lysates samples